

ARMY RESEARCH LABORATORY



Spectroscopic Studies of Gas-Phase  
ArCH(D) Van der Waals (vdW)  
Complexes: II. Observation and Analysis  
of B-X Electronic Transitions of ArCD

Rosario C. Sausa  
Michael J. McQuaid  
George W. Lemire

ARL-TR-1159

August 1996

DTIC QUALITY INSPECTED 4

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION IS UNLIMITED.

19961008 019

## **NOTICES**

Destroy this report when it is no longer needed. DO NOT return it to the originator.

Additional copies of this report may be obtained from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Road, Springfield, VA 22161.

The findings of this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

The use of trade names or manufacturers' names in this report does not constitute indorsement of any commercial product.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE August 1996		3. REPORT TYPE AND DATES COVERED Final, Jul 92-Jul 93
4. TITLE AND SUBTITLE Spectroscopic Studies of Gas-Phase ArCH(D) Van der Waals (vdW) Complexes: II. Observation and Analysis of B-X Electronic Transitions of ArCD			5. FUNDING NUMBERS PR: 1L162618AH80	
6. AUTHOR(S) Rosario C. Sausa, Michael J. McQuaid, and George W. Lemire*				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory ATTN: AMSRL-WT-PC Aberdeen Proving Ground, MD 21005-5066			8. PERFORMING ORGANIZATION REPORT NUMBER ARL-TR-1159	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES *National Research Council (NRC)/U.S. Army Research Laboratory (ARL) Postdoctoral Research Associate; present address is Joint Contact Directorate, U.S. Army Dugway Proving Ground, Dugway, UT 84022-5000.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The ArCD van der Waals (vdW) complex has been generated in the gas phase by the 248-nm photolysis of CDBr <sub>3</sub> entrained in a supersonic jet of argon. Rotationally resolved spectra associated with the complex have been observed by laser-induced fluorescence in the vicinity of the CD B <sup>2</sup> Σ <sup>-</sup> - X <sup>2</sup> Π (1,0) band near 366 nm. Bands at 27310.7 cm <sup>-1</sup> and 27327.8 cm <sup>-1</sup> are rotationally analyzed using computer simulations and assigned as transitions to the B state (1,0 <sup>0</sup> ,0) and (1,1 <sup>0</sup> ,0) vibrational levels, respectively. The vibronic structure of the ArCD excited states has also been analyzed using a model based on hindered internal rotation. The results are compared to those reported for ArCH.				
14. SUBJECT TERMS ArCH, ArCD, potential energy surfaces			15. NUMBER OF PAGES 17	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

INTENTIONALLY LEFT BLANK.

## ACKNOWLEDGMENTS

We wish to thank Professor M. Heaven, Emory University, and Dr. A. Kotlar, U.S. Army Research Laboratory (ARL), for many valuable discussions. We are also grateful to Professor P. Dagdigian, Johns Hopkins University, and Drs. N. Garland and H. Nelson, U.S. Naval Research Laboratory, for providing us with the deuterated compounds. This work was supported by the ARL Combustion Research and Environmental Research Programs. Purchase of equipment through the Productivity Capital Investment Program (R. Sausa) and support from the National Research Council (NRC)/ARL Postdoctoral Program (G. Lemire) are gratefully acknowledged.

INTENTIONALLY LEFT BLANK.

## TABLE OF CONTENTS

	<u>Page</u>
ACKNOWLEDGMENTS .....	iii
LIST OF FIGURES .....	vii
1. INTRODUCTION .....	1
2. EXPERIMENTAL .....	1
3. RESULTS/DISCUSSION .....	2
4. SUMMARY/CONCLUSION .....	7
5. REFERENCES .....	11
DISTRIBUTION LIST .....	13

INTENTIONALLY LEFT BLANK.



## LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1. Laser-induced fluorescence scan of ArCD excited near the CD $B^2\Sigma^- - ^2\Pi$ (1,0) band .	3
2. Laser-induced excitation spectrum and simulations (---) of the feature labeled A in Figure 1 in the region of the CD $B^2\Sigma^- - ^2\Pi$ (1,0) transitions . . . . .	5
3. A plot of the $V_{2,0}$ and $V_{2,2}$ coefficients of the renormalized spherical harmonic expansion used to model the intermolecular potential . . . . .	8
4. Angular dependence of the ArCH(D) (X,A'') and (X,A') potential energy surfaces corresponding to $V_{2,0} = 20 \text{ cm}^{-1}$ , $V_{2,2} = -28 \text{ cm}^{-1}$ . . . . .	9

INTENTIONALLY LEFT BLANK.

## 1. INTRODUCTION

In recent years, there has been a considerable interest in characterizing van der Waals (vdW) complexes formed by a diatomic radical and a rare-gas atom. Representative studies of such systems are given in references [1–15]. Complexes involving the CH radical are of particular interest. The radical plays an important role in atmospheric and interstellar chemistry and mediates a wide variety of reactive processes in combustion and flames. Numerical modeling of these various environments depends on the measurement and/or theoretical calculation of numerous bimolecular reaction rates involving many different collisional partners. A prerequisite for deriving reaction rates is a representation of the intermolecular potential surface on which the interaction takes place. However, generating a potential function that adequately represents the surface can be quite difficult, particularly for reactive collisions. Spectroscopic investigations of rare gas - CH(D) systems are a step in this direction.

In a recent study, we reported the observation of laser excitation spectra associated with gas-phase ArCH where the electronic states of the complex correlate with the  $^1S_0$  state of Ar and the  $X^2\Pi_r$  and  $B^2\Sigma^-$  states of CH [2]. It was found that the ground state geometry was T-shaped, while the B state was linear. The ground state geometry was inferred based on the similarity between the observed rotational energy level structure and that of bent triatomic molecules. The geometry of the excited state was based on a tentative vibronic assignment and a modeling of the energy separation between the proposed bending levels assuming a hindered internal rotation description [14]. Here, we report the observation and analysis of rotationally resolved spectra of the ArCD vdW complex, excited in the region of the CD  $B^2\Sigma^- - X^2\Pi$  (1,0) band. The results are compared to those obtained for the ArCH complex.

## 2. EXPERIMENTAL

The molecular beam apparatus used to obtain the spectra reported in this report has been described previously [3]. The ArCD complexes were formed via a supersonic free jet expansion.  $CDBr_3$  vapor was entrained in Ar gas and expanded into a vacuum chamber through an extender channel mounted on a pulsed valve (Jordan Co.). The backing pressure for the expansion was nominally 100–120 psi. CD radicals were generated in the early stages of the expansion by photolyzing the  $CDBr_3$ /Ar mixture with the focused output of an excimer laser (Lambda Physik, EMG 150 MSC) operating at 248 nm. (Comparable ArCD signals were obtained using  $CDBr_2Cl$  as a precursor.) Typical pulse energy at the photolysis region was approximately 10 mJ. Laser-induced fluorescence was used to probe the ArCD and

CD products in the region of the CD  $B^2\Sigma^- - X^2\Pi$  (1,0) band near 366 nm. The probe beam was directed normal to the photolysis laser beam and molecular beam a few millimeters from the extender channel. Tunable laser radiation with a nominal pulse energy of 5 mJ was generated using a XeCl excimer pumped dye laser (Lumonics Hyper EX-400 and Hyper DYE-300) operating with PBD dye. In this wavelength region, the line width of the probe laser is approximately  $0.09\text{ cm}^{-1}$ . The ArCD complex was also probed in the vicinity of the CD  $B^2\Sigma^- - X^2\Pi$  (0,0) transition near 388 nm, but poor signal-to-noise (S/N) levels precluded an analysis of the rotational energy level structure. Under similar experimental conditions, the S/N ratio for the ArCD complexes probed near the CD (0,0) and (1,0) bands was a factor of 5–10 weaker than that for the ArCH complexes probed near the CH (0,0) and (1,0) bands. The fluorescence from the ArCD transitions excited in the region of the CD  $B^2\Sigma^- - X^2\Pi$  (1,0) band was monitored near the CD  $B^2\Sigma^- - X^2\Pi$  (1,1) transition centered at 401 nm. The fluorescence was band-pass filtered by a 0.25-m monochromator (McPherson, Model 218) and detected with a photomultiplier tube (EMI-9789-QA). After amplification, the output signal was directed to a gated integrator (Stanford Research Systems) with a PC interface for data acquisition. The spectra were calibrated with a wavemeter (Candela, Model LS-1) whose calibration was crosschecked by measuring the  $P_1(1)$  and  $R_2(1)$  CH rotational frequencies and comparing them to previously established values [16–17]. Absolute wavenumbers could be determined to  $\pm 0.7\text{ cm}^{-1}$ , and wavenumber differences could be measured to  $0.09\text{ cm}^{-1}$ .

### 3. RESULTS/DISCUSSION

Figure 1 presents a survey fluorescence excitation spectrum obtained near the CD  $B^2\Sigma^- - X^2\Pi$  (1,0) band. This spectrum has not been corrected for variation in laser intensity or detection sensitivity. The sharp features at low energies are CD B-X (1,0) rotational lines. To the blue of these transitions is a sequence of broad, red-shaded features (A–H) associated with the ArCD complex. A broad continuum is also observed in this region and is presumably associated with bound-free transitions of the complex. At higher resolution, partially resolved rovibronic transitions are observed for features A and B, but not for features C–H.

Good empirical fits of bands A and B were obtained using a model based on  $\Sigma - \Sigma$  type transitions [2]. In this model, the term values for the ground state rotational energy levels are calculated using the expressions

$$F_1''(N) = B''N(N+1) + 0.5\gamma''N \quad (1a)$$




Figure 1. Laser-induced fluorescence scan of ArCD excited near the CD  $B^2\Sigma^- - ^2\Pi (1,0)$  band.

and

$$F_2''(N) = B''N(N+1) - 0.5\gamma''(N+1), \quad (1b)$$

where  $N$  is an integer quantum number and  $B$  is the effective rotational constant. The terms containing  $\gamma''$  were included to allow for Coriolis coupling related splitting in the ground state. Justification for these functional forms is provided in Lemire et al. [2]. Following the treatment by Chang et al. [9], the excited state term values are given by

$$F_1'(N) = B'N(N+1) + 0.5\mu N \quad (2a)$$

and

$$F_2'(N) = B'N(N+1) - 0.5\mu(N+1). \quad (2b)$$

In equations 2a and 2b,  $\mu$  may be further expanded as

$$\mu = (\kappa p^2 / (N(N+1))) + \gamma', \quad (3)$$

where  $\kappa$  is the coefficient for the spin-rotation term associated with vibronic levels and  $p$  is the projection of the vibrational angular momentum on the axis of the complex. The selection rule  $\Delta N = \pm 1$  leads to six branches, which we label  $P_1$ ,  $P_2$ ,  $P_{12}$ ,  $R_1$ ,  $R_2$ , and  $R_{12}$ .

Features A and B were recorded at an effective resolution of  $0.09 \text{ cm}^{-1}$ . In contrast to the ArCH B-X spectra, where the R and P branches were clearly observed at this resolution, lower S/N levels prevent a straightforward assignment of these features. Rather, the initial assignment was based on correspondence with the results obtained for ArCH. This procedure is justified because, considered as a pseudodiatom, the reduced masses ( $\mu$ ) of ArCD and ArCH are similar ( $\mu_{\text{Ar-CD}} = 10.0$  vs.  $\mu_{\text{Ar-CH}} = 9.8$ ). Thus, the rotational energy level structure and rotational constants of these complexes should be comparable. This is in contrast to the case of CD vs. CH, where the reduced mass of CD ( $\mu_{\text{CD}} = 1.71$ ) is much larger than that of CH ( $\mu_{\text{CH}} = 0.92$ ), and, consequently, the two have much different rotational energy level separations.

To verify assignments, obtain rotational constants, and determine the band origins, graphical simulations were generated using a multivariable computer program based on the energy level expressions and selection rules given previously, and  $\Sigma$ - $\Sigma$  line strengths. Besides the spectroscopic constants of the ground and excited states, the program variables included the laser line width and the population of the ground state based on a Boltzmann distribution. It was found that the correspondence with ArCH bands was not sufficient to confidently assign the rotational spectra. Thus, it was further assumed that both A and B originated from the same ground state. This assumption is based on the fact that the complexes were generated in a free jet expansion with a temperature of approximately 10 K.

Figure 2 presents a comparison of the experimentally obtained spectrum for feature A and two simulations, labeled  $S_1$  and  $S_2$ , based on different rotational assignments. Both simulations reasonably reproduce the band. The simulation labeled  $S_1$  is based on  $B'' = 0.172 \pm 0.004 \text{ cm}^{-1}$ ,  $B' = 0.107 \pm 0.004 \text{ cm}^{-1}$ , and  $\gamma'' \leq (\pm)0.019 \text{ cm}^{-1}$ , while  $S_2$  is based on  $B'' = 0.168 \pm 0.004 \text{ cm}^{-1}$ ,  $B' = 0.102 \pm 0.004 \text{ cm}^{-1}$ , and  $\gamma'' = (\pm)0.035 \text{ cm}^{-1}$ . Both simulations yield a band origin value of  $27310.7 \text{ cm}^{-1}$ .

To differentiate between  $S_1$  and  $S_2$ , we attempted to simulate feature B using the two sets of ( $B''$ ,  $\gamma''$ ) values obtained from the analysis of feature A. A better fit of the feature was obtained using the ( $B''$ ,  $\gamma''$ ) values employed in  $S_1$  ( $B'' = 0.172 \pm 0.004 \text{ cm}^{-1}$ ,  $\gamma = 0.019 \text{ cm}^{-1}$ ). However, this fit did not reproduce several lines observed from  $27,323$  through  $27,327 \text{ cm}^{-1}$ . These "extra" lines may be associated




Figure 2. Laser-induced excitation spectrum and simulations (---) of the feature labeled A in Figure 1 in the region of the CD  $B^2\Sigma^- - {}^2\Pi(1,0)$  transitions. (Simulations  $S_1$  and  $S_2$  are based on two different rotational assignments.  $S_1$  yields rotational constant values of  $B'' = 0.172 \pm 0.004 \text{ cm}^{-1}$ ,  $B' = 0.107 \pm 0.004 \text{ cm}^{-1}$ , and  $\gamma'' = 0.195 \text{ cm}^{-1}$ ; while  $S_2$  yields values of  $B'' = 0.168 \pm 0.004 \text{ cm}^{-1}$ ,  $B' = 0.102 \pm 0.004 \text{ cm}^{-1}$ , and  $\gamma''$ . Both simulations yield a band origin value of  $27310.7 \text{ cm}^{-1}$ .)

with other products formed in the jet or another ArCD band, so we cannot rule out the assignment based on  $S_1$  on this basis alone. Using  $B'' = 0.172 \text{ cm}^{-1}$  and  $\gamma'' = 0.019 \text{ cm}^{-1}$ , the best simulation of feature B yields a value of  $B' = 0.112 \pm 0.004 \text{ cm}^{-1}$  for the excited state rotational constant and a value of  $27327.8 \text{ cm}^{-1}$  for the band origin. The ground state  $B''$  values obtained from the analysis of features A and B correspond to an average Ar-CD center of mass separation ( $R_0''$ ) of  $3.13 \text{ \AA}$ , which is similar to the  $R_0''$  value of  $3.11 \text{ \AA}$  reported for the ground state of the ArCH complex. The  $R_0'$  values fall in the range of those reported for ArCH,  $3.97\text{--}4.61 \text{ \AA}$ , and indicate that the Ar-CD vdW bond is lengthened upon electronic excitation.

Using the notation  $(v_1, r^p, v_s)$ , where  $v_1$  is the quanta of CD stretch,  $r$  is the quanta of bend/hindered internal rotation of CD with respect to Ar,  $p$  is the projection of the vibrational angular momentum on the intermolecular axis of the complex, and  $v_s$  is the quanta of CD-Ar stretch, the features labeled A and B are assigned as  $(1,0^0,0)$  and  $(1,1^p,0)$ , respectively. These assignments are based on a comparison with the results obtained for ArCH. One observation is considered important in this regard: the small but definite increase in the  $B'$  value for the higher energy transition. For ArCH, the  $B'$  value for the level assigned as  $(0,1^0,0)$  was also slightly larger than the  $B'$  value for the  $(0,0^0,0)$  level ( $0.114$  v.  $0.111$   $\text{cm}^{-1}$ ). These results would be realized in the case where the D/H atom points away from the argon atom in the vibrationless state, and introducing the bending/hindered internal rotation reduces the Ar-CD center-of-mass separation. For feature B,  $p$  could not be determined from the rotational analysis. Spectral simulation in which  $\kappa$  was varied over a considerable range ( $-2B'' \leq \kappa \leq 2B''$ ) did not qualitatively affect the fits.

In order to gain further insight into the nature of the ground state intermolecular potential, and aid in the rotational assignment of feature A, we have analyzed the spectroscopic data in terms of a general model based on hindered internal rotation which neglects bend-stretch coupling. The model was developed by Dubernet, Flower, and Hutson [14] and later expanded by Chang et al. [9] to include the Coriolis coupling interaction. The intermolecular potential of the complex is modeled by

$$V_{inter} = \sum_{l,m} V_{l,m}(R) C_m^l(\theta,0) = V_{1,0} C_0^1(\theta,0) + V_{2,0} C_0^2(\theta,0) + V_{2,2} C_2^2(\theta,0) + \dots \quad (4)$$

where  $\theta$  is the angle between the monomer and complex axis, and  $V_{l,m}(R)$  are coefficients of the renormalized spherical harmonic expansion terms  $[C_m^l(\theta,0)]$ . These coefficients are varied to reproduce the energy level structure separations observed in the spectra. For the following discussion, we restrict ourselves to expansion terms with  $l \leq 2$ .

For the ArCH(D) ground state, the geometry of the intermolecular potential is reflected in the Coriolis splitting of the rotational levels. Each end-over-end rotational state ( $N$ ) correlates with a pair of degenerate, parity doublets  $N = J \pm 1/2$ . The degeneracy of this state is lifted for  $V_{l,m}$  not equal to zero. The energy levels for a  $V_{l,m}$  combination were calculated by diagonalizing the effective Hamiltonian derived by Chang et al. [9]. The functional form of the ground state energy levels, equations (1a) and (1b), is particular to the case of a potential dominated by  $V_{2,m}$  terms and with  $V_{1,0} = 0$ . For ArCH,  $\gamma''$  was observed to be  $(\pm)0.021$   $\text{cm}^{-1}$ . Assuming that the barriers to hindered internal rotation for the ArCH



and ArCD complexes are similar, there do not exist expansion coefficients which yield  $\gamma''_{\text{ArCH}} = (\pm)0.021 \text{ cm}^{-1}$  and  $\gamma''_{\text{ArCD}} \leq 0.034 \text{ cm}^{-1}$ . Thus, the  $\gamma (\leq 0.019 \text{ cm}^{-1})$  obtained from the simulation labeled  $S_1$  is not consistent with equal barriers for ArCH and ArCD in this description. The  $\gamma (0.035 \text{ cm}^{-1})$  obtained from simulation  $S_2$  is consistent with this expectation. Plots of the solutions for  $\gamma_{\text{ArCH}} = (\pm)0.021 \text{ cm}^{-1}$  and  $\gamma_{\text{ArCD}} = (\pm)0.035 \text{ cm}^{-1}$  are presented in Figure 3. In this figure, the solutions for each case form a closed curve in the  $V_{2,0}$ - $V_{2,2}$  plane. Agreement between these sets of solutions is only observed at  $V_{2,0} = 20 \text{ cm}^{-1}$ ,  $V_{2,2} = -28 \text{ cm}^{-1}$ , and  $V_{2,0} = -28 \text{ cm}^{-1}$ ,  $V_{2,2} = 28 \text{ cm}^{-1}$ . The uncertainties in  $V_{2,0}$  and  $V_{2,2}$  are estimated to be approximately  $\pm 7 \text{ cm}^{-1}$  and  $\pm 3 \text{ cm}^{-1}$ , respectively. It is interesting to note that the solution corresponds to  $V_{l,m}$  combination in each set for which  $|V_{2,2}|$  is a maximum. Dagdigian et al. [18] have shown that  $V_{22}$  is negative for the case of a  $\text{CH } ^2\Pi$  monomer in a  $\pi^1$  configuration inelastically scattered from a rare gas atom. Thus, the solution  $V_{2,0} = 20 \text{ cm}^{-1}$ ,  $V_{2,2} = -28 \text{ cm}^{-1}$  describes the intermolecular potential if such an analysis is warranted. Figure 4 shows the ground state  $A''$  and  $A'$  potential energy surfaces corresponding to this solution. While it is recognized that the uncertainty in assigning the rotational structure of feature A leaves room for doubt about this description, the T-shaped geometry for the  $A''$  state and the lack of a barrier to internal rotation for the  $A'$  surface are in agreement with the results obtained from multireference configuration interaction calculations reported by Alexander et al. [19].

#### 4. SUMMARY/CONCLUSION

The ArCD vdW complex has been detected near the CD  $B^2\Sigma^- - ^2\Pi (1,0)$  band by laser-induced fluorescence. Rotationally resolved spectra of the complex have been recorded and analyzed by computer simulations and a model based on hindered internal rotation. The analysis, coupled with the results from spectroscopic and numerical studies of ArCH, indicates that the ground state interaction between  $\text{CH(D)} (X^2\Pi)$  and  $\text{Ar}(^1S_0)$  has  $A''$  symmetry and has a (global) minimum associated with a (nearly) T-shaped configuration. We anticipate that these results will provide a useful foundation/starting point for future efforts to characterize the potential energy surfaces for reactive collisional (combustion) systems involving the CH radical.




Figure 3. A plot of the  $V_{2,0}$  and  $V_{2,2}$  coefficients of the renormalized spherical harmonic expansion used to model intermolecular potential.




Figure 4. Angular dependence of the ArCH(D) (X,A'') and (X,A') potential energy surfaces corresponding to  $V_{2,0} = 20 \text{ cm}^{-1}$ ,  $V_{2,2} = -28 \text{ cm}^{-1}$ .

INTENTIONALLY LEFT BLANK.

## 5. REFERENCES

1. Heaven, M. C. "Spectroscopy and Dynamics of Open-Shell van der Waals Molecules." Ann. Rev. Phys. Chem., vol. 43, p. 283, 1992.
2. Lemire, G. W., M. J. McQuaid, A. J. Kotlar, and R. C. Sausa. "Investigation of the Gas-Phase B-X Electronic Spectra of CH-Ar by Laser-Induced Fluorescence." Journal of Chemical Physics, vol. 99, p. 91, 1993.
3. Lemire, G. W., and R. C. Sausa. "Detection and Spectroscopic Studies of Gas-Phase Hydroxyl-Krypton by Laser-Induced Fluorescence." Journal of Physical Chemistry, vol. 96, p. 4821, 1992.
4. Green, W. H., Jr., and M. I. Lester. "A Perturbation Theory Guide to Open-Shell Complexes: OH-AR ( $X^2\Pi$ )." Journal of Chemical Physics, vol. 96, p. 2573, 1992.
5. Randall, R. W., C.-C. Chang, and M. I. Lester. "Electronic Spectroscopy of NH ( $c^1\Pi$ ,  $a^1\Delta$ ) + Ar Complexes." Chemical Physics Letters, vol. 200, p. 113, 1992.
6. Chang, B.-C., L. Lu, D. Cullin, B. Rehfuss, J. Williamson, T. A. Miller, W. M. Fawzy, X. Zheng, S. Fei, and M. C. Heaven. "Rotational, Fine and Hyperfine Structure in the High Resolution Electronic Spectrum of ArOH and ArOD." Journal of Chemical Physics, vol. 95, p. 7086, 1991.
7. Yang, M.-C., A. P. Salzberg, B.-C. Chang, C. C. Carter, and T. A. Miller. "Electronic Spectroscopy and Excited State Dynamics of the ArSH Complex." Journal of Chemical Physics, vol. 98, p. 4301, 1993.
8. Miller, J. C., and W.-C. Cheng. "Multiphoton Ionization of NO-Rare Gas van der Waals Species." Journal of Physical Chemistry, vol. 89, p. 1647, 1989.
9. Chang, B.-C., J. Cullin, J. Williamson, J. Dunlop, B. Rehfuss, and T. Miller. "High Resolution Electronic Spectroscopy of NeOH." Journal of Chemical Physics, vol. 96, p. 3476, 1992.
10. Miller, J. C. "The  $A^2\Sigma^+$  State of Rare Gas - NO van der Waals Molecules Probed by 1+1 Multiphoton Ionization Spectroscopy." Journal of Chemical Physics, vol. 90, p. 4031, 1989.
11. Alexander, M. H. "Rotationally Inelastic Collisions Between a Diatomic Molecule in a  $^2\Pi$  Electronic State and a Structureless Target." Journal of Chemical Physics, vol. 76, p. 5974, 1982.
12. Fawzy, W. M., and J. T. Hougen. "Rotational Energy Levels and Line Intensities for  $^{2S+1}\Lambda - ^{2S+1}\Lambda$  and  $^{2S+1}(\Lambda\pm 1) - ^{2S+1}\Lambda$  Transitions in Diatomic Molecule van der Waals Bonded to a Closed Shell Partner." Journal of Molecular Spectroscopy, vol. 137, p. 154, 1989.
13. Chakravarty, C., D. C. Clary, A. Degli Esposti, and H.-J. Werner. "Calculations on Vibrational Predissociation of Ar-OH ( $A^2\Sigma^+$ )." Journal of Chemical Physics, vol. 95, p. 8149, 1991.
14. Dubernet, M.-L., D. Flower, and J. Hutson. "The Dynamics of Open-Shell van der Waals Complexes." Journal of Chemical Physics, vol. 94, p. 7602, 1991.

15. Dubernet, M.-L., P. A. Tuckey, and J. M. Hutson. "Parity Doubling in Open Shell van der Waals Complexes." Chemical Physics Letters, vol. 193, p. 355, 1992.
16. Gerö, L. "Das Bandenspectrum des CD Molekuls." Z. Phys., vol. 117, p. 709, 1941.
17. Para, A. "Analysis of the  $B^2\Sigma^- - X^2\Pi$  (0,0) Band in the  $^{12}\text{C}^1\text{H}$ ,  $^{12}\text{C}^2\text{H}$ , and  $^{13}\text{C}^1\text{H}$  Isotopic Radicals." J. Phys. B: At. Mol. Opt., vol. 24, p. 3179, 1991.
18. Dagdigan, P. J., M. H. Alexander, and K. Liu. "The Inelastic Scattering of  $^2\Pi$  [Case (b)] Molecules and an Understanding of the Differing Doublet Propensities for Molecules of  $\pi$  vs.  $\pi^3$  Orbital Occupancy." Journal of Chemical Physics, vol. 91, p. 839, 1989.
19. Alexander, M. H., S. Gregurick, P. J. Dagdigan, G. W. Lemire, M. J. McQuaid, and R. C. Sausa. "Potential Energy Surfaces for the Interaction of CH ( $X^2\Pi$ ,  $B^2E^-$ ) with Ar and an Assignment of the Stretch-Bend Levels of the ArCH(B) van der Waals Molecule." Journal of Chemical Physics, vol. 101, p. 4547, 1994.

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
2	DEFENSE TECHNICAL INFO CTR ATTN DTIC DDA 8725 JOHN J KINGMAN RD STE 0944 FT BELVOIR VA 22060-6218

1	DIRECTOR US ARMY RESEARCH LAB ATTN AMSRL OP SD TA 2800 POWDER MILL RD ADELPHI MD 20783-1145
---	---

3	DIRECTOR US ARMY RESEARCH LAB ATTN AMSRL OP SD TL 2800 POWDER MILL RD ADELPHI MD 20783-1145
---	---

1	DIRECTOR US ARMY RESEARCH LAB ATTN AMSRL OP SD TP 2800 POWDER MILL RD ADELPHI MD 20783-1145
---	---

ABERDEEN PROVING GROUND

2	DIR USARL ATTN AMSRL OP AP L (305)
---	---------------------------------------

<u>NO. OF COPIES</u>	<u>ORGANIZATION</u>
2	JOINT CONTACT POINT DIR USA DUGWAY PROVING GRND ATTN G LEMIRE DUGWAY UT 84022-5000
19	DIR, USARL ATTN: AMSRL-WT-P, A. HORST AMSRL-WT-PA, T. MINOR M. MCQUAID (10 CPS) R. SAUSA (5 CPS) AMSRL-WT-PC, R. FIFER K. MCNESBY



## USER EVALUATION SHEET/CHANGE OF ADDRESS

This Laboratory undertakes a continuing effort to improve the quality of the reports it publishes. Your comments/answers to the items/questions below will aid us in our efforts.

1. ARL Report Number/Author ARL-TR-1159 (Sausa) Date of Report August 1996
2. Date Report Received \_\_\_\_\_
3. Does this report satisfy a need? (Comment on purpose, related project, or other area of interest for which the report will be used.) \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_
4. Specifically, how is the report being used? (Information source, design data, procedure, source of ideas, etc.) \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_
5. Has the information in this report led to any quantitative savings as far as man-hours or dollars saved, operating costs avoided, or efficiencies achieved, etc? If so, please elaborate. \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_
6. General Comments. What do you think should be changed to improve future reports? (Indicate changes to organization, technical content, format, etc.) \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

CURRENT  
ADDRESS

\_\_\_\_\_  
Organization

\_\_\_\_\_  
Name

\_\_\_\_\_  
Street or P.O. Box No.

\_\_\_\_\_  
City, State, Zip Code

7. If indicating a Change of Address or Address Correction, please provide the Current or Correct address above and the Old or Incorrect address below.

OLD  
ADDRESS

\_\_\_\_\_  
Organization

\_\_\_\_\_  
Name

\_\_\_\_\_  
Street or P.O. Box No.

\_\_\_\_\_  
City, State, Zip Code

(Remove this sheet, fold as indicated, tape closed, and mail.)  
(DO NOT STAPLE)

---

**DEPARTMENT OF THE ARMY**


**OFFICIAL BUSINESS**

**BUSINESS REPLY MAIL**

**FIRST CLASS PERMIT NO 0001,APG,MD**

**POSTAGE WILL BE PAID BY ADDRESSEE**

**DIRECTOR  
U.S. ARMY RESEARCH LABORATORY  
ATTN: AMSRL-WT-PC  
ABERDEEN PROVING GROUND, MD 21005-5066**



**NO POSTAGE  
NECESSARY  
IF MAILED  
IN THE  
UNITED STATES**

